

## Composition:-

1 tablet of: Fluconazole...	150 mg
1 tablet of: Azithromycin dihydrate	
Equivalent to Azithromycin base...	1000 mg
2 tablets each of Secnidazole...	1000 mg

## Therapeutic Indications & Usage:-

**FLUZAN-AS** is indicated for treatment of mixed vaginal infections or as empirical cure of suspected mixed vaginal infections such as vulvovaginitis, bacterial vaginosis and trichomoniasis; it is also indicated in syndromic management of pelvic inflammatory disease.

## Contraindications:-

**FLUZAN-AS** is contraindicated in patients with known hypersensitivity to this product or any of its ingredients, or to nitroimidazoles or macrolides. Co-administration of terfenadine and cisapride with Fluconazole (in **FLUZAN-AS**) is contraindicated.

**FLUZAN-AS** is not advocated during pregnancy, lactation and in children.

## Precautions:-

**FLUZAN-AS** is to be advocated with due precaution in hepatic and renal disease, and potentially proarrhythmic conditions in view potential of Fluconazole and Azithromycin to induce liver and kidney dysfunction, and cardiac conduction abnormalities.

**FLUZAN-AS** should be advocated with care in diarrhea since Azithromycin usage could be associated with pseudo membranous colitis.

## Adverse Reactions:-

Single dose of 150 mg Fluconazole (in **FLUZAN-AS**) can cause mild to moderate headache, nausea, abdominal pain, diarrhea, dyspepsia, dizziness and taste perversion. Other rare side effects due to Fluconazole (in **FLUZAN-AS**) include vomiting, seizures, cardiac conduction abnormalities, skin lesions, hematopoietic disturbances, electrolyte and lipid changes.

Azithromycin (in **FLUZAN-AS**) can cause gastrointestinal side effects such as nausea, vomiting, diarrhea and abdominal pain. Other adverse reactions due to Azithromycin (in **FLUZAN-AS**) include fatigue, asthenia, paresthesia, malaise, hypersensitivity and skin eruptions. Cholestatic jaundice, palpitations, chest pain and arrhythmias, genitourinary affections such as monilia, vaginitis, nephritis and acute renal failure, nervous system manifestations like dizziness, headache, vertigo, somnolence, hyperactivity, nervousness, agitation, syncope, aggressive reaction and anxiety, as well as hearing disturbances and taste perversion can occur with Azithromycin (in **FLUZAN-AS**). Azithromycin (in **FLUZAN-AS**), like other antimicrobials, can cause pseudo membranous colitis.

Secnidazole (in **FLUZAN-AS**) can cause mild side effects of gastrointestinal disturbances, and dizziness.

Rarely, angioedema and anaphylactic reactions can occur with **FLUZAN-AS** ingredients.

## Drug Interactions:-

Fluconazole (in **FLUZAN-AS**) can interact with oral hypoglycemics, coumarins, phenytoin, cyclosporine, rifampicin, theophylline, astemizole, rifabutin, tacrolimus, terfenadine, cisapride and short-acting benzodiazepines.

Azithromycin (in **FLUZAN-AS**) absorption can be reduced by antacids having magnesium or aluminium. Elevation of serum levels of digoxin, ergot derivatives, Triazolam and drugs metabolized by liver cytochrome P450 can occur with Azithromycin (in **FLUZAN-AS**).

Secnidazole (in **FLUZAN-AS**) can potentiate the action of anticoagulants.

## Recommended Dosage & Administration:-

**ADULTS:** 1 **FLUZAN-AS** Combikit as single oral dose, or as prescribed. Both the partners should preferably be treated simultaneously.

## Packaging Information:-

**FLUZAN-AS Combikit:** Blisters of 4s.

1 tablet of: Fluconazole... 150 mg

1 tablet of: Azithromycin dihydrate Equivalent to Azithromycin base...1000 mg

2 tablets each of Secnidazole...1000 mg

Trusted Quality Products From GCR



**Green Cross<sup>®</sup>**  
**Remedies**

Trusted Quality Products From GCR